

# Section 3: Environmental Control

## 3.2 Environmental Decontamination



Jointly prepared by  
Infection Control Branch, Centre for Health Protection, Department of Health

and

Central Committee on Infectious Diseases, Hospital Authority



## **Acknowledgements:**

**We would like to thank all individuals who have contributed to the contents of this infection control guideline:**

**Special acknowledgement: This guideline has been reviewed by**

**Dr Raymond LAI, COS (MIC), Prince of Wales Hospital, Hospital Authority**

**Dr. WL Lim, Head, Public Health Laboratory Services Branch, Centre for Health Protection, Department of Health**

## **Experts of external consultation parties**

Members of Central Committee on Infectious Diseases, Hospital Authority  
Chairman, Scientific Committee on Infection control, Centre for health protection  
Chairman, Infection Control Committee, Department of Health  
Infection control Nursing Sub-Committee of COC (N), Hospital Authority  
Representatives of Privates Hospitals

## **Members of Guideline Development work group**

Dr. Wong, Ada, Medical Officer, Elderly Health Services, DH (September 2005 to March 2006)  
Dr. Yip, Lisa, Medical Officer, Elderly Health Services, DH  
Ms. Chow, Sin Cheung, ICN, Ruttonjee and Tang Shiu Kin Hospital  
Ms. Fong, Oi Wah, Nursing Officer, Public Health Service Branch, CHP  
Ms. Ho, Yuk Yin, ICN, Tung Wah Hospital  
Mr. Kan, Chun Hoi, SNO, Tuen Mun Hospital  
Ms. Kwok, Stella, ICN, Hong Kong Buddhist Hospital  
Ms. Lau, Joan, Nursing Officer, Elderly Health Services, DH (from 25 October 2005)  
Ms. Leung, Fung Yee, DOM, Princess Margaret Hospital  
Mr. Leung, Tsz Kin, ICN, Prince of Wales Hospital  
Ms. Luk, Amy, SNO, Hong Kong Baptist Hospital  
Ms. Sit, Amy, ICN, Tai Po Hospital  
Ms. Tam, Oi Yi, Catherine, ICN, Pamela Youde Nethersole Eastern Hospital  
Mr. Tsoi, Wai Lun, ICN, United Christian Hospital  
Ms. Wong, Susanna, Nursing Officer, Elderly Health Services, DH (September to October 2005)  
Ms. Woun, Babbitt, ICN, Tuen Mun Hospital  
Mr. Yu, Man Kit, ICN, Queen Elizabeth Hospital

## **Core-working group members of Infection Disease Control Training Centre, HAHO / Infection Control Branch, Centre for Health Protection:**

Dr. Yung Wai Hung, Raymond, Head  
Dr. Wong, Tin Yau, Associate Consultant  
Dr. Chan, Kai Ming, Associate Consultant  
Dr. Chuang, Vivien, Associate Consultant  
Dr. Leung, Yiu Hong, Medical Officer (August 2005 to March 2006)  
Dr. Luk, Shik, Medical Officer (Jul 2006 to Dec 2006)  
Dr. Lam, Bosco, Medical Officer  
Mr. Lee, Kai Yip Ralph, Occupational Hygienist  
Ms. Chan, Toi Lan, Nursing Officer  
Ms. Chan, Wai Fong, Advanced Practice Nurse (Nov 2004 to December 2005)  
Ms. Leung, Suk Yee Jane, Advanced Practice Nurse  
Ms. Chan, Mei Mei, Registered Nurse  
Ms. Kwok, Man Kee, Registered Nurse (August 2005 to September 2005)  
Ms. Ng, Wai Po, Registered Nurse (July 2005 to August 2005)  
Ms Wong, Kwan Wai, Registered Nurse (March 2006 to Jan 2007)  
Ms. Yuen, Woon Wah, Registered Nurse

## Contents

1. [Category of Environmental Surfaces](#)
2. [Decontamination of Environmental Surfaces](#)
3. [Handling Spills of Blood, Body Fluids, or other Potentially Infectious Materials](#)
4. [Environmental Control in Special Areas](#)
  - 4.1. Neonatal unit
  - 4.2. Areas with immunocompromised patients
  - 4.3. Operating room

## [Table 1: Common Disinfectants Used for Environmental Cleaning in Hospitals](#)

## [References](#)

Microbiologically contaminated inanimate surfaces generally are not directly associated with transmission of infections to human beings (1). The main concern is still transmission indirectly via hand contact with the contaminated surface. Cleaning the environmental surfaces is fundamental in reducing their potential contribution to infections (2-4).

### **1 Category of Environmental Surfaces**

Environmental surfaces can be further divided into medical equipment surfaces (e.g. haemodialysis machines, x-ray machines, instrument carts and dental units) and housekeeping surfaces (e.g. floors, walls, tabletops, bedside rails) (1).

### **2 Decontamination of Environmental Surfaces**

- 2.1 To facilitate the environmental cleaning, belongings of patients/ residents are preferably stored inside the wardrobe, locker or in a covered container.
- 2.2 Environmental surfaces should be cleansed by “active damp scrubbing” (5) to avoid creation of aerosols or splashing, or dust dispersion (1-3, 6-8).

- 2.3 Environmental surfaces should be cleaned regularly, for example, daily, and decontaminated promptly when visibly soiled (1-2, 7). A general rule of environmental cleaning is to start cleaning at the highest point and work towards the lowest; and from outside to the inside (8).
- 2.4 High-touch surfaces should be cleansed more frequently, for example, at least daily, than minimal touch surfaces (1,5).
- 2.5 In general, detergent is adequate for cleaning the environmental (housekeeping and equipment) surfaces in healthcare facilities and community, including institutions and households (1-3, 9-14).
- 2.6 Cleaning and disinfection procedures in HA hospitals should be carried out in accordance with local policy.
- 2.7 Areas with frequent invasive procedures, such as haemodialysis unit or blood donation centre, low level disinfection, for example, 500ppm sodium hypochlorite solution (one part of 5.25% sodium hypochlorite solution add in 99 parts of water) may be employed for decontaminating environmental surfaces, because minute spillage may be difficult to recognize (10-11,15). When visible blood spills occur, decontaminating procedure stated as Point 3 should be followed.
- 2.8 In the areas accommodating patients with organisms that may contaminate the environment easily, for example in outbreak situation, environmental surfaces may be cleansed and disinfected with a higher level of disinfection strategy, for example, intermediate-level disinfection using one part of household bleach (5.25% sodium hypochlorite solution) in 49 parts of water (about 1,000 ppm available chlorine) at least daily (1-2, 5, 16-18). Use 70% alcohol solution to disinfect equipment's metal surface.

- 2.9 For environmental surfaces which are contaminated by uncommon infective materials, for example, Creutzfeldt-Jakob disease (CJD), please refer to *fact sheet on Infection Control of Creutzfeldt-Jakob Disease (CJD)*, prepared by Central Committee on Infectious Diseases of Hospital Authority:  
[http://www3.ha.org.hk/idtc/document/Infection\\_Control\\_for\\_CJD.pdf](http://www3.ha.org.hk/idtc/document/Infection_Control_for_CJD.pdf)

### **3 Handling Spills of Blood, Body Fluids, or other Potentially Infectious Materials**

Other potentially infectious materials means (19):

- Human body fluids include semen, vaginal secretions, cerebrospinal fluid, synovial fluid, pleural fluid, pericardial fluid, peritoneal fluid, amniotic fluid, saliva in dental procedures, any body fluid that is visibly contaminated with blood, and all body fluids in situations where it is difficult or impossible to differentiate between body fluids, such as mixing of faeces and blood;
- Any unfixed tissue or organ (other than intact skin) from a human (living or dead)

3.1 If spills occur, the surfaces should be decontaminated immediately (1, 10, 19-22) with adequate staff protection (1, 10). For recommendation of appropriate protection, please refer to Section Personal Protective Equipment of ICB Infection Control Guidelines.

3.2 For spillage of blood and other potentially infectious materials, cleanse the visible matter with disposable absorbent material. Mop the area with a cloth or paper towels wetted with one part of household bleach (5.25% hypochlorite solution) in 4 parts of water, leave for 10 minutes. Then rinse with water.

Small spill of blood can also be removed by applying chlorine-releasing granules or powder directly to the spill. This can then be removed using one or more paper towels or wipes (23).

3.3 For spillage of other body fluids such as vomitus, cleanse the visible matter with disposable absorbent material. Mop the area with a cloth or paper towels with one part of household bleach (5.25% hypochlorite solution) in 49 parts of water, leave for 15-30 minutes. Then rinse with water.

#### 4 Environmental Control in Special Areas

##### 4.1 Neonatal unit

When performing cleansing procedure with disinfectant to environmental surfaces of neonatal area, avoid unnecessary exposure of neonates to disinfectant residue, for example, do not use disinfectant to cleanse incubators during the infant's stay, and rinse the disinfectant-treated surfaces with water (1).

##### 4.2 Areas with immunocompromised patients

Areas of immunocompromised patients, such as haematopoietic stem cell transplant recipients, wet-mopping method should be used to clean the surfaces (1, 24).

The doors of immunocompromised patients' rooms should be closed when vacuuming, waxing, or buffing corridor floors to minimize exposure to dust (1).

##### 4.3 Operating room

Wet vacuum or mop the operating room floor with disinfectant after the last operation of the day or night (25).

When the surfaces or equipment is visibly soiled or contaminated with blood or other body fluids, clean and disinfect the affected areas before the next operation (25).

**Section 3: Environmental Control**  
**3.2 Environmental Decontamination**



Issue Date: 9 August 2007 (Advanced Draft)

Page 5 of 10

---

Tacky mats at the entrance of operating room or individual operating room for infection control purpose is not necessary (1, 22).



Jointly prepared by  
Infection Control Branch, Centre for Health Protection, Department of Health  
and  
Central Committee on Infectious Diseases, Hospital Authority



**Table 1: Common Disinfectants Used for Environmental Cleaning in Hospitals**

Disinfectants	Recommended Use	Precautions
Sodium hypochlorite 1% in-use dilution, one part of household bleach (5.25% hypochlorite solution) in 4 parts of water	Disinfection of material contaminated with blood and other potentially infectious materials.	<ul style="list-style-type: none"> <li>✧ Should be used in well ventilated areas</li> <li>✧ Protective clothing required while handling and using undiluted disinfectants</li> <li>✧ Do not mix with strong acids to avoid release of chlorine gas</li> <li>✧ Corrosive to metals</li> </ul>
Alcohol (70%) Isopropyl, ethyl alcohol, methylated spirit.	Smooth metal surfaces, tabletops and other surfaces on which bleach cannot be used.	<ul style="list-style-type: none"> <li>✧ Flammable, toxic, to be used in well-ventilated area, avoid inhalation.</li> <li>✧ Kept away from heat source, electrical equipment, flames, hot surfaces.</li> <li>✧ Allow it to dry completely, particularly when using diathermy as it can cause diathermy burns.</li> </ul>

**Note: A neutral detergent and water solution should be used for all routine and general cleaning. When a disinfectant is required for surface cleaning, e.g. after spillage or contamination with blood or body fluids, the manufacturer's recommendations for use and occupational health and safety instructions should be followed.**

## References

1. Centers for Disease and Prevention. Guidelines for environmental infection control in health-care facilities. MMWR 2003;52 (RR-10):1-48.
2. World Health Organization-West Pacific & South-East Asia Regions. Practical guidelines for infection control in health care facilities. [online] 2004 [cited 2005 Aug 24]. Available from URL: [http://www.searo.who.int/LinkFiles/Publications\\_PracticalguidelinSEAROpub-41.pdf](http://www.searo.who.int/LinkFiles/Publications_PracticalguidelinSEAROpub-41.pdf)
3. World Health Organization-West Pacific Region. Interim guidelines for national SARS preparedness. [online] 2003 [cited 2005 Feb 22]. Available from: URL: <http://www.wpro.who.int/sars/docs/interimguidelines/default.asp>
4. World Health Organization. Healthy villages. A guide for communities and community health workers. [online] 2002 [cited 2005 Aug 5] Available from: URL: <http://whqlibdoc.who.int/publications/2002/9241545534.pdf>
5. Muto CA, Jernigan A, Ostrowsky BE, Richet HM, Jarvis WR, Boyce JM, et al. SHEA guideline for preventing nosocomial transmission of multidrug-resistant strains of staphylococcus aureus and enterococcus. Infect Control Hosp Epidemiol 2003;24(5):362-86.
6. Smith TL, Iwan PC, Olson SB, Rupp ME. Environmental contamination with vancomycin-resistant enterococci in an outpatient setting. Infect Control Hosp Epidemiol 1998;19(7):515-8.
7. Griffith CJ, Cooper RA, Gilmore J, Davies C, Lewis M. An evaluation of hospital cleaning regimes and standards. J Hosp Infect 2000; 45(1):19-28.

8. National Health Service. Healthcare facilities cleaning manual. [online] 2004 [cited 2005 Feb 23]. Available from: URL: [http://patientexperience.nhsestates.gov.uk/clean\\_hospitals/ch\\_content/cleaning\\_manual/cleaning\\_methods.asp](http://patientexperience.nhsestates.gov.uk/clean_hospitals/ch_content/cleaning_manual/cleaning_methods.asp)
9. Centers for Disease Control and Prevention. Guideline for handwashing and hospital environmental control. MMWR 1988;37(24):1-25.
10. Rutala WA. APIC guideline for selection and use of disinfectants. Am J Infect Control 1996;24(4):313-42.
11. Rutala WA, Weber DJ. Disinfection and sterilization in healthcare facilities: draft CDC guidelines. In: Rutala WA, editor. Disinfection, sterilization and antisepsis. Principles, practices, challenges, and new research. Washington: APIC; 2004. p.42-75.
12. Allerberger F, Ayliffe G, Bassetti M, Braveny I, Bucher A, Damani N, et al. Routine surface disinfection in health care facilities: should we do it? Am J Infect Control 2002;30(5):318-9.
13. Rutala WA, Weber DJ. Selection and use of disinfectants in healthcare. In: Mayhall, CG, editor. Hospital epidemiology and infection control. 3<sup>rd</sup> ed. Philadelphia: Lippincott Williams & Wilkins; 2004. p.1473-522.
14. Dharan S, Mourouga P, Copin P, Bessmer G, Tschanz B, Pittet D. Routine disinfection of patients' environmental surfaces. Myth or reality? J Hosp Infect 1999;42(2):113-7.
15. Centers for Disease Control and Prevention. Recommendations for preventing transmission of infections among chronic hemodialysis patients. MMWR 2001;50(RR-5),1-46.

16. Rampling A, Wiseman S, Davis L, Hyett AP, Walbridge AN, Payne GC, et al. Evidence that hospital hygiene is important in the control of methicillin resistant staphylococcus aureus. *J Hosp Infect* 2001;49(2):109-16.
17. Rutala WA, Weber DJ. Uses of inorganic hypochlorite (Bleach) in health-care facilities. *Clin Microbiol Rev* 1997;10(4):597-610.
18. Lemmen SW, Hafner H, Zolldann D, Stanzel S, Lütticken R. Distribution of multi-resistant Gram-negative versus gram-positive bacteria in the hospital inanimate environment. *J Hosp Infect* 2004;56(3):191-7.
19. US Department of Labor, Occupational Safety and Health Administration. Bloodborne pathogens: definitions: other potentially infectious materials (29CFR 1910.1030b). [online] 1991 [cited 2007 May 9]. Available from URL: [http://www.osha.gov/pls/oshaweb/owadisp.show\\_document?p\\_table=STANDARDS&p\\_id=10051](http://www.osha.gov/pls/oshaweb/owadisp.show_document?p_table=STANDARDS&p_id=10051)
20. US Department of Labor, Occupational Safety and Health Administration. Bloodborne pathogens: housekeeping (29CFR 1910.1030 d4iiA). [online] 1991 [cited 2007 May 9]. Available from URL: [http://www.osha.gov/pls/oshaweb/owadisp.show\\_document?p\\_table=STANDARDS&p\\_id=10051](http://www.osha.gov/pls/oshaweb/owadisp.show_document?p_table=STANDARDS&p_id=10051)
21. Wearmouth P. (2004). Cleanliness matters in NHS. [online] 2004 [cited 2005 Feb 23]. Available from: URL: <http://www.nursing-standard.co.uk/archives/ns/vol18-44/pdfs/v18n44NScleanliness.pdf>
22. Hospital Authority Central Committee on Infectious Disease. Guideline on transport of clinical specimens and infectious substances. [online] 2004 [cited 2006 Dec 18].

**Section 3: Environmental Control**  
**3.2 Environmental Decontamination**



Issue Date: 9 August 2007 (Advanced Draft)

Page 10 of 10

---

Available from: URL:  
<http://ha.home/ho/ps/Guidelinesontransportofclinicalspecimens.pdf>

23. Jean L, Dee M. Infection control in the community. NY: Churchill Livingstone; 2003.
24. Centers for Disease Control and Prevention. Guidelines for preventing opportunistic infections among haematopoietic stem cell transplant recipients. MMWR 2000;49 (RR-10):1-125.
25. Mangram AJ, Horan TC, Pearson ML, Silver LC, Jarvis WR, The Hospital Infection Control Practices Advisory Committee. Guideline for prevention of surgical site infection, 1999. Infect Control Hosp Epidemiol 1999;20(4):247-78.



Jointly prepared by  
Infection Control Branch, Centre for Health Protection, Department of Health  
and  
Central Committee on Infectious Diseases, Hospital Authority

