

# FLU EXPRESS



*Flu Express* is a weekly report produced by the Respiratory Disease Office of the Centre for Health Protection. It monitors and summarizes the latest local and global influenza activities.

## Local Situation of Influenza Activity (as of Mar 30, 2016)

**Reporting period: Mar 20 – 26, 2016 (Week 13)**

- Some surveillance data have shown that the local influenza activity has continued to decrease from the peak level but still remained elevated. It is expected that the influenza activity will remain elevated for some time. The public should continue to be vigilant.
- The Centre for Health Protection (CHP) has collaborated with the Hospital Authority (HA) and private hospitals to reactivate the enhanced surveillance for severe seasonal influenza cases (i.e. influenza-associated admissions to intensive care unit (ICU) or deaths) among patients aged 18 or above since Jan 29, 2016. As of Mar 30, 284 adult severe cases (including 120 deaths) were recorded. Separately, 17 cases of severe paediatric influenza-associated complication/death among patients aged below 18 years (including one death) were recorded during the period.
- Influenza can cause serious illnesses in high-risk individuals and even healthy persons. Given that seasonal influenza vaccines are safe and effective, all persons aged 6 months or above except those with known contraindications are recommended to receive influenza vaccine for personal protection.
- Eligible children (aged between six months and less than 6 years, or 6 years old or above attending a kindergarten or child care centre in Hong Kong), elderly (aged 65 years or above) and eligible persons with intellectual disabilities can be subsidised for seasonal influenza vaccination from enrolled private doctors participating in the Government's vaccination subsidy schemes starting from Oct 15, 2015. Elderly aged 65 or above living in the community can also receive free vaccination from General Out-patient Clinics under the HA and designated Elderly Health Centres of the Department of Health since Nov 10, 2015. Details are available from the vaccination schemes website ([http://www.chp.gov.hk/en/view\\_content/17980.html](http://www.chp.gov.hk/en/view_content/17980.html)).

## Influenza-like-illness surveillance among sentinel general outpatient clinics and sentinel private doctors, 2012-16

In week 13, the average consultation rate for influenza-like illness (ILI) among sentinel general outpatient clinics (GOPCs) was 9.0 ILI cases per 1,000 consultations, which was higher than 8.2 recorded in the previous week (Figure 1, left). The average consultation rate for ILI among sentinel private doctors was 59.9 ILI cases per 1,000 consultations, which was lower than 69.7 recorded in the previous week (Figure 1, right).




Figure 1 ILI consultation rate at sentinel GOPCs (left) and private doctors (right), 2012-16

## Laboratory surveillance, 2012-16

Among the respiratory specimens received in week 13, 679 (20.95%) were tested positive for seasonal influenza viruses, including 271 (8.36%) influenza A(H1), 17 (0.52%) influenza A(H3), 363 (11.20%) influenza B and 28 (0.86%) influenza C. The percentage of respiratory specimens tested positive for seasonal influenza viruses last week was 20.95%, which was higher than 18.46% recorded in the previous week (Figure 2). Among the influenza viruses detected in the last week, the proportions of B, A(H1), C and A(H3) were 53.5%, 39.9%, 4.1% and 2.5% respectively. The proportion of influenza B among positive influenza detections has been increasing steadily and has overtaken A(H1) to become the most commonly detected subtype in last three weeks.




Figure 2 Percentage of respiratory specimens tested positive for influenza viruses, 2012-16 (upper: overall positive percentage, lower: positive percentage by subtypes)

## Influenza-like illness outbreak surveillance, 2012-16

In week 13, 24 ILI outbreaks occurring in schools/institutions (affecting 115 persons) were recorded, as compared to 38 outbreaks (affecting 199 persons) recorded in the previous week (Figure 3). In the first 4 days of week 14 (Mar 27 to 30, 2016), 4 institutional ILI outbreaks (affecting 20 persons) were recorded.




Figure 3 ILI outbreaks in schools/institutions, 2012-16

## Rate of influenza-like illness syndrome group in accident and emergency departments, 2012-16<sup>#</sup>

In week 13, the rate of the influenza-like illness syndrome group in the accident and emergency departments (AED) was 246.7 (per 1,000 coded cases), which was similar to the rate of 244.9 in the previous week (Figure 4).

*#Note: The influenza-like illness syndrome group includes codes such as influenza, upper respiratory tract infection, fever, cough, throat pain, and pneumonia.*




Figure 4 Rate of ILI syndrome group in AED, 2012-16

## Influenza associated hospital admission rates and deaths in public hospitals based on discharge coding, 2009-16

In week 13, the admission rates in public hospitals with principal diagnosis of influenza for persons aged 0-4 years, 5-9 years, 10-64 years and 65 years or above were 2.74, 1.01, 0.15 and 0.58 cases (per 10,000 people in the age group) respectively, as compared to 2.99, 1.45, 0.14 and 0.57 cases in the previous week (Figure 5). Weekly number of deaths with any diagnosis of influenza is also shown in Figure 5.




Figure 5 Influenza associated hospital admission rates and deaths, 2009-16

## Fever surveillance at sentinel child care centres/ kindergartens, 2012-16

In week 13, 1.06% of children in the sentinel child care centres/ kindergartens (CCC/ KG) had fever ( $38^{\circ}\text{C}$  or above) as compared to 1.14% in the previous week (Figure 6).




Figure 6 Percentage of children with fever at sentinel CCC/ KG, 2012-16

## Fever surveillance at sentinel residential care homes for the elderly, 2012-16

In week 13, 0.13% of residents in the sentinel residential care homes for the elderly (RCHes) had fever ( $38^{\circ}\text{C}$  or above) as compared to 0.10% in the previous week (Figure 7).




Figure 7 Percentage of residents with fever at sentinel RCHE, 2012-16

## Influenza-like illness surveillance among sentinel Chinese medicine practitioners, 2012-16

In week 13, the average consultation rate for ILI among Chinese medicine practitioners (CMPs) was 1.62 ILI cases per 1,000 consultations as compared to 2.83 in the previous week (Figure 8).




Figure 8 ILI consultation rate at sentinel CMP, 2012-16

## Surveillance of severe influenza cases

(Note: The data reported are provisional figures and subject to further revision)

- Since activation of the enhanced surveillance for severe influenza infection on Jan 29, 2016, a total of 284 adult severe cases (including 120 deaths) and 17 paediatric severe cases (including one death) were recorded (as of Mar 30)(Figure 9). Among them, 203 patients had infection with influenza A(H1N1)pdm09, 71 patients with influenza B, 15 patients with influenza A(H3N2), one patient with influenza C, 10 patients with influenza A pending subtype and one patient with both influenza A(H3N2) and B. In the last winter season in early 2015, 647 adult cases (including 501 deaths) and 18 paediatric cases (including 1 death) were filed.

### Enhanced surveillance for severe seasonal influenza (Aged 18 years or above)

- In week 13, 30 cases of influenza associated ICU admission/death were recorded, in which 13 of them were fatal. In the first 4 days of week 14 (Mar 27 to 30), 17 cases of influenza associated ICU admission/death were recorded, in which 6 of them were fatal.

### Surveillance of severe paediatric influenza-associated complication/death (Aged below 18 years)

- In week 13, there were two cases of severe paediatric influenza-associated complication. In the first 4 days of week 14 (Mar 27 to 30), there were two cases of severe paediatric influenza-associated complication. The details are as follow:

| Reporting week | Age       | Sex  | Complication              | Influenza subtype      |
|----------------|-----------|------|---------------------------|------------------------|
| 13             | 23 months | Male | Pneumonia and myocarditis | Influenza A(H1N1)pdm09 |
| 13             | 13 years  | Male | Sepsis                    | Influenza B            |
| 14             | 10 years  | Male | Encephalopathy            | Influenza B            |
| 14             | 5 months  | Male | Pneumonia                 | Influenza B            |




Figure 9 Weekly number of severe influenza cases recorded during influenza seasons, 2012-2016

Remark: The surveillance system for severe influenza cases aged 18 years or above was only activated intermittently during influenza seasons.

## Surveillance of oseltamivir resistant influenza A(H1N1)pdm09 virus infection

- In week 13 and the first 4 days of week 14 (Mar 27 to 30, 2016), there were no new reports of oseltamivir (Tamiflu) resistant influenza A(H1N1)pdm09 virus infection. There are totally 47 reports of oseltamivir resistant influenza A(H1N1)pdm09 virus detected in Hong Kong since 2009.

## Global Situation of Influenza Activity

- In the United States (week ending Mar 19, 2016), influenza activity decreased slightly but remained elevated. The proportion of outpatient visits for ILI was 3.2%, which was above the national baseline of 2.1%.
- In Canada (week ending Mar 19, 2016), the influenza activity remained near its peak, but has begun to decrease. Influenza A(H1N1)pdm09 is the most common circulating virus subtype.
- In the United Kingdom (week ending Mar 19, 2016), the influenza indicators remained stable. Influenza B circulation further increased. The weekly ILI consultation rate has increased and was above the medium intensity threshold in England. The ILI rates in Scotland decreased slightly. A slight increase in the ILI rate was noted in Northern Ireland, and that in Wales remained similar to previous week. The percentage of positive influenza detection increased to 25.7%, which was above the threshold for 2015/16 season of 7.4%.
- In Europe (week ending Mar 20, 2016), influenza was still widespread in countries in the western part of the Region, but majority of countries reported decreasing or stable trends. There was a shift towards the circulation of influenza B virus, which constituted 66% of influenza virus detections in sentinel samples.
- In Mainland China (week ending Mar 20, 2016), the influenza activities in both southern and northern China were at seasonal epidemic levels and continued to increase. Influenza outbreaks further increased. Proportion of influenza B detection overtook that of influenza A.
- In Taiwan (week ending Mar 19, 2016), the overall influenza activity continued to decrease. The numbers of ILI consultations in AED and severe influenza cases have been decreasing. Influenza B constituted about 60% of influenza detection. During the past four weeks, the antigenic match between the seasonal influenza vaccine and the circulating influenza virus strains were 100% in H1N1 and H3N2 viruses, but 43% in influenza B virus.
- In Japan (week ending Mar 20, 2016), the influenza season has started since early January. The average number of reported ILI cases per sentinel site decreased to 21.13 in the week ending Mar 20 from 28.20 in the previous week, but still much higher than the baseline level of 1.00.

### Sources:

Information have been extracted from the following sources when updates are available: [United States Centers for Disease Control and Prevention](#), [Public Health Agency of Canada](#), [Public Health England](#), [Joint European Centre for Disease Control and Prevention-World Health Organization/Flu News Europe](#), [Chinese National influenza Center](#), [Taiwan Centers for Disease Control](#) and [Japan Ministry of Health](#).