



**衛生防護中心**  
Centre for Health Protection

## **Scientific Committee on AIDS and STI (SCAS)**

### **Using Antiretrovirals for Post Exposure Prophylaxis against HIV in the Non-occupational Setting – Position Statement of the SCAS**

#### **Preamble**

Human immunodeficiency virus (HIV) is transmitted by three major routes: blood borne, sexual, and mother-to-child. Early attempts with pharmacologic agents to prevent HIV infection after exposure were in the occupational setting and empiric in nature.

2. Since 1997, the use of HIV post exposure prophylaxis in the occupational setting (PEP) has been made the standard of care in Hong Kong as well as overseas, following guidelines by the then Scientific Committee on AIDS.<sup>i</sup> In overseas countries, post exposure prophylaxis with antiretrovirals in the non-occupational setting (nPEP) has also been used, mainly targeted against exposure from sexual contact and shared injecting equipment. Although a number of guidelines on nPEP are now available in the US and some countries of Europe,<sup>ii,iii,iv,v</sup> they are noted for their variability in the antiretroviral recommendations and the circumstances where they should be used. This has caused confusion to both the public and health care providers.

3. The SCAS deliberated on this issue in Nov 2005 by examining the available scientific evidence, reviewing reported experience in local and overseas providers, and considering its ethical implications. This position statement represents the consensus of the Committee regarding nPEP.



衛生防護中心乃衛生署  
轄下執行疾病預防  
及控制的專業架構

*The Centre for Health  
Protection is a*

*professional arm of the  
Department of Health for  
disease prevention and  
control*

## Limited scientific basis of nPEP

4. Proof-of-concept studies with simian immunodeficiency virus (SIV) and HIV-2 in nonhuman primates showed moderate degrees of success if treatment was started very early, preferably at the time of exposure.<sup>vi,vii,viii,ix</sup> Treatment initiated more than 72 hours after exposure would very likely be useless. This essentially imposed a 72-hour window of intervention.

5. In one retrospective study, zidovudine initiated within 48 h of birth apparently was able to decrease the transmission risk from 26.6% to 9.3%.<sup>x</sup> Human data have otherwise been largely based on a case-control study in health care workers (HCW), in which use of zidovudine was associated with an 81% reduction in transmission risk.<sup>xi</sup> The quality of data however was compromised by the small number of cases ( $n=31$ ) and controls ( $n=679$ ) originating from different cohorts. This and other methodological issues with the study may have overestimated the efficacy.<sup>xii</sup> Nevertheless these data formed the basis of one major cost effectiveness study on nPEP.<sup>xiii</sup> This showed that the intervention would be cost-effective only in the event that the source was known to be HIV infected and the exposure was receptive anal sex or shared injecting equipment. Tempering optimism with nPEP is that failures have occurred.<sup>xiv</sup>

6. *All in all, scientific research to date has not matured to the point of recommending nPEP as standard of care. Any use of nPEP would therefore be exceptional and should be considered only in the event of high-risk exposure to a source known to be HIV positive. If 72 hours have elapsed since exposure, nPEP should not be prescribed. Were nPEP to be given, the recipient should fully understand the toxicity and experimental nature of this intervention.*

## Antiretroviral regimens

7. Highly active antiretroviral therapy (HAART) generally comprises at least three drugs from two or more classes. Its potency and durability have been proven in established HIV infection, having significantly lowered morbidity and mortality in Hong Kong.<sup>xv,xvi</sup>

8. Nevertheless, the studies on which PEP was based largely employed monotherapy. It is rational to the notion that combination regimen, especially HAART, may be more efficacious than monotherapy. Using HAART in prevention against mother-to-child transmission has lowered the risk to less than 2%,<sup>xvii</sup> levels never achieved in any clinical trial using just one drug. In addition, the new generation of antiretrovirals has notably a less severe profile of adverse effects, making the number of drugs *per se* not as important in predicting toxicity in a recipient. In designing a regimen, the toxicity profile should be one that has the least impact on the recipient. For example, efavirenz should be withheld from women of childbearing age because of potential

teratogenicity. Nevirapine should not be prescribed for nPEP at all because of pronounced liver toxicity in HIV negative persons.<sup>xviii</sup>

9. Although certain antiretrovirals have been suggested to be preferable based on inference from their mechanisms of action or in vitro effects, it is unclear if any possesses enhanced prophylactic efficacy. There being no clinical comparative data, one therefore cannot argue convincingly in favour of any particular drug to be included for the purpose of efficacy. Rather, the possibility of drug resistance should be considered and factored in during the selection of drugs. Worldwide, primary drug resistance is emerging as a significant problem, especially in some developed countries where antiretrovirals are widely prescribed. Secondary drug resistance is more common, however, and is usually secondary to poor drug adherence. If available, the treatment history and resistance test results in the source patient are helpful in determining the appropriate antiretroviral regimen. Conceivably, poor adherence to nPEP may impact on the viral resistance profile should transmission occur. It is therefore important to monitor adherence closely and manage toxicity expectantly and appropriately.

10. *Thus members of SCAS are of the view that HAART is preferred were nPEP to be given. Its composition is dictated by the toxicity profile and the possibility of drug resistance. To prevent the emergence of drug resistance, nPEP should be given in settings supervised by physicians experienced in antiretroviral therapy.*

### **Other aspects of managing non-occupational HIV exposure**

11. Exposure to HIV by the sexual route or injecting drug use differs from that of occupational setting in several important aspects. Risk assessment may be difficult because of the often imprecise recount of the sexual act or drug taking episode. Clients present relatively late. Exposure source is not readily available for testing. There are other sexually transmitted infections or blood-borne pathogens which pose significant, if not more, threats. As exposure is behaviour-mediated, there is also a high likelihood of repetition.

12. *nPEP is but one facet of the overall management of non-occupational exposure to HIV. In fact, over-emphasis on nPEP by the client or the health care provider risks overlooking the importance of risk-reduction counselling on safer sex and safe injection practice. Most STIs are of higher transmissibility than HIV and should be screened for after sexual exposure. Depending on circumstances, blood-borne pathogens such as hepatitis B and C should also be managed by serology testing and immunisation where applicable. Tetanus vaccination history should also be reviewed after percutaneous exposure.*

13. A frontline health care provider may not be equipped to handle

all aspects of management and referral to specialists is advised after immediate management.

Centre for Health Protection  
March 2006

The copyright of this paper belongs to the Centre for Health Protection, Department of Health, Hong Kong Special Administrative Region. Contents of the paper may be freely quoted for educational, training and non-commercial uses provided that acknowledgement be made to the Centre for Health Protection, Department of Health, Hong Kong Special Administrative Region. No part of this paper may be used, modified or reproduced for purposes other than those stated above without prior permission obtained from the Centre.

## References

- i Scientific Committee on AIDS and Scientific Working Group on Viral Hepatitis Prevention. Procedure for management of needlestick injury or mucosal contact with blood or body fluids – general guidelines for hepatitis B, C and HIV prevention. Sep 1997 (Available at [http://www.info.gov.hk/aids/english/publications/pubsearch\\_1.htm#HIV%20Management%20and%20Clinical%20Guidelines](http://www.info.gov.hk/aids/english/publications/pubsearch_1.htm#HIV%20Management%20and%20Clinical%20Guidelines). Accessed 3 Jan 2006)
- ii New York State Department of Health. HIV prophylaxis following non-occupational exposure including sexual assault. Dec 2004 (Available at [http://www.hivguidelines.org/public\\_html/npep/npep.pdf](http://www.hivguidelines.org/public_html/npep/npep.pdf). Accessed 3 Jan 2006)
- iii The California Task force on Non-Occupational PEP and the California Department of Health Services, Office of AIDS. Offering HIV postexposure prophylaxis (PEP) following non-occupational exposures. Recommendations for health care providers in the state of California. June 2004 (Available at <http://www.aids-ed.org/pdf/guidelines/pepassault.pdf>. Accessed Nov 30, 2005)
- iv US CDC. Antiretroviral postexposure prophylaxis after sexual, injection-drug use, or other nonoccupational exposure to HIV in the United States. MMWR 2005;24(RR-2)
- v Almeda J, Casabona J, Simon B, et al on behalf of the Euro-NONOPEP Project Group. Proposed recommendations for the management of HIV post-exposure prophylaxis after sexual, injecting drug or other exposures in Europe. Eurosurveillance 2004;9(2):35-43
- vi Spira AI, Marx PA, Patterson BK, et al. Cellular targets of infection and route of viral dissemination after an intravaginal inoculation of simian immunodeficiency virus into rhesus macaques. J Exp Med 1996;183:215-25
- vii Tsai CC, Follis KE, Sabo A, et al. Prevention of SIV infection in macaques by (R)-9-(2-phosphonmethoxyprophyl)adenine. Science 1995;273:1197-9
- viii Bottiger D, Johansson N-G, Samuelsson B, et al. Prevention of simian immunodeficiency virus, SIVsm, or HIV-2 infection in cynomolgus monkeys by pre- and postexposure administration of BEA-005. AIDS 1997;11:157-62
- ix Tsai CC, Emau P, Follis KE, et al. Effectiveness of postinoculation (R)-9-(2-phosphonmethoxyprophyl)adenine treatment for prevention of persistent simian immunodeficiency virus SIVmne infection depends critically on timing of initiation and duration of treatment. J Virol 1998;72:4265-73
- x Wade NA, Birkhead GS, Warren BL, et al. Abbreviated regimens of zidovudine prophylaxis and perinatal transmission of human immunodeficiency virus type 1. N Engl J Med 1998;339:1409-14
- xi Cardo DM, Culver DH, Ciesielski CA, et al. A case-control study of HIV seroconversion in health care workers after percutaneous exposure. N Eng J Med 1997;337:1485-90
- xii Henderson DK. Postexposure treatment of HIV – taking some risks for safety’s sake [editorial]. N Engl J Med 1997;337:1542-3
- xiii Pinkerton SD, Holtgrave DR, Bloom FR. Cost-effectiveness of post-exposure prophylaxis following sexual exposure to HIV. AIDS 1998;12:1067-78
- xiv Rolance ME, Neilands TB, Krone MR, et al. Seroconversion following nonoccupational postexposure prophylaxis against HIV. Clin Infect Dis 2005;41:1507-13.
- xv Chan KCW, Cheng LS, Chan WK, Wong KH. Highly active antiretroviral

- 
- therapy per se decreased mortality and morbidity of advanced human immunodeficiency virus disease in Hong Kong. *Chin Med J* 2005;118(16):1338-45
- xvi Wong KH, Chan KCW, Lee SS. Delayed progression to death and to AIDS in a Hong Kong cohort of patients with advanced HIV type 1 disease during the era of highly active antiretroviral therapy. *Clin Infect Dis* 2004;39:853-60
- xvii McGowan JP, Crand M, Wiznia AA, et al. Combination antiretroviral therapy in human immunodeficiency virus-infected pregnant women. *Obstet Gynecol* 1999;94:641-6
- xviii US CDC. Serious adverse events attributed to nevirapine regimens for postexposure prophylaxis after HIV exposure – worldwide, 1997-2000. *MMWR* 2001;49:1153-7