

Surveillance And Epidemiology Branch

本署檔號	Our Ref.	:	(19) in DH SEB CD/8/91/1
來函檔號	Your Ref	:	
電 話	Tel.	:	
傳 真	Fax No.	:	

監測及流行病學處

30 April 2009

Dear Principal / Person-in-charge,

Update on Swine Influenza

Further to my letter on 27 April 2009, I would like to update you on the latest situation of swine influenza (H1N1).

On 29 April 2009, the World Health Organization raised the **level of influenza pandemic alert from phase 4 to phase 5**, which is a strong signal that a pandemic is imminent. As of 8:30 am today, affected areas with cases of confirmed human swine influenza include Mexico, the United States, Canada, Spain, the United Kingdom, New Zealand, Israel, Austria and Germany.

I would like to take this opportunity to remind kindergartens and schools to take all precautions against swine influenza as mentioned in our previous letter. Please also remind your students to observe the health advice and precautionary measures on the attached swine influenza pamphlet and avoid all non-essential travel to human swine flu affected areas, particularly Mexico.

The Department of Health has set up a 24-hour hotline (2125 1111) to facilitate enquiries from people who developed flu-like symptoms and had travelled to human swine flu affected areas during the 7-day period before onset of symptoms. People who presented with fever, cough, sore throat or have difficulty in breathing and had travelled to the affected areas during the 7-day period before onset of symptoms, they should call the hotline immediately.

If you notice an increase in respiratory illnesses or absenteeism among your staff or students, please report to our Central Notification Office (Tel: 2477-2772; Fax: 2477-2770) for immediate epidemiological investigations and outbreak control.

衛生防護中心乃衛生署 轄下執行疾病預防 及控制的專業架構 The Centre for Health Protection is a professional arm of the Department of Health for disease prevention and control

_ _ _ _ _ _ _ _ _ _ _ _ _ _ .

For the latest information and health education material on human swine influenza, please see our CHP website at www.chp.gov.hk.

Yours sincerely,

per

(Dr SK CHUANG) Consultant Community Medicine (Communicable Disease) Surveillance and Epidemiology Branch Centre for Health Protection

Human Swine Influenza

Background

A number of confirmed human cases of swine influenza A/H1N1 (swine flu) infection have been reported in Mexico and several other countries.

Originally known to circulate among pig populations, the swine flu viruses do occasionally infect human beings. In this present international swine flu outbreak, human-to-human transmission has occurred.

Symptoms

The symptoms of human swine influenza are usually similar to those of human seasonal influenza and include fever, lethargy, loss of appetite and coughing. Some people infected with swine flu may also have a runny nose, sore throat, nausea, vomiting and diarrhoea.

Mode of transmission

Human-to-human transmission of swine flu is thought to occur in the same way as seasonal flu is spread among people, mainly through coughing or sneezing. People may also become infected by touching objects soiled with flu viruses and then touching their nose or mouth.

Swine flu has not been shown to be transmissible to people through eating properly handled and cooked pork or pork products. Cooking pork to an internal temperature of 70° C (160° F) kills the swine flu virus.

Management

People who develop flu symptoms should put on a mask and consult a doctor as soon as possible. Those who have been to affected places or been exposed to sick persons should tell the doctor the travel and contact history. Antiviral agents can reduce the severity and duration of illness but must be used under doctor's prescription. It is important for people not to self-medicate.

Prevention

Since H1N1 swine flu viruses are very different from human H1N1 viruses, vaccines for human seasonal flu will not provide protection against H1N1 swine flu viruses.

Members of the public should observe the following precautionary measures:

- Keep hands clean and wash hands properly. Alcohol-based handrub is also effective when hands are not visibly soiled.
- Avoid touching mouth, nose or eyes.
- Wash hands with liquid soap promptly if they are dirtied by respiratory secretions, e.g. after sneezing or coughing.
- Cover nose and mouth when sneezing or coughing.
- Do not spit. Always wrap nasal and mouth discharges with tissue paper, and dispose of the tissue paper properly in a rubbish bin with a lid.
- Wear a mask when respiratory symptoms or fever develop. See a doctor right away.
- Do not go to work or school if you develop influenza-like symptoms.

Avoid travelling to affected places unless it is absolutely necessary. If such a visit is unavoidable:

- During the trip: put on a mask and do not have contact with sick people.
- After returning: pay close attention to your health and wear masks for 7 days. Seek medical consultation from public clinics or hospitals right away if fever or influenza-like symptoms appear.

Printed in April 2009



www.chp.gov.hk 2833 0111

