

COVID-19 & FLU EXPRESS



COVID-19 & Flu Express is a weekly report produced by Surveillance Division of the Communicable Disease Branch of the Centre for Health Protection. It monitors and summarises the latest local and global COVID-19 and influenza activities.

Local Situation of COVID-19 Activity (as of Sep 2, 2026)

Reporting period: Aug 23, 2026 – Aug 29, 2026 (Week 35)

- The latest surveillance data showed that the overall local activity of COVID-19 continued to decline.
- The Centre for Health Protection (CHP) has been closely monitoring the local prevalence of SARS-CoV-2 variants. The latest sewage surveillance data and genetic analysis of positive respiratory specimens showed that PQ.16.1.1 is the most prevalent variant in Hong Kong. PQ.16.1.1 is a descendant lineage of NB.1.8.1, which is a descendant of JN.1. The World Health Organization (WHO) listed PQ.16.1.1 as a variant under monitoring (VUM) on Jul 27, 2026. The WHO stated that existing COVID-19 vaccines are expected to be effective against PQ.16.1.1, and there is no evidence to suggest that PQ.16.1.1 will cause more serious diseases.
- Members of the public are advised to maintain strict personal and environmental hygiene at all times for personal protection against COVID-19 infection and prevention of the spread of the disease in the community. High-risk people (e.g. persons with underlying medical conditions or persons who are immunocompromised) should adopt additional measures to protect themselves such as wearing mask properly when going to public places. For other details, please visit the COVID-19 information page (<https://www.chp.gov.hk/en/healthtopics/content/24/102466.html>).
- Members of the public are advised to take note of the latest recommendations on the use of COVID-19 vaccines in Hong Kong to protect themselves from serious outcomes of COVID-19. High-risk priority groups are recommended to receive a dose of COVID-19 vaccine at least six months since the last dose or infection, regardless of the number of doses received previously. For more details, please visit (https://www.chp.gov.hk/files/pdf/consensus_recommendations_on_the_use_of_covid-19_vaccines_in_hong_kong_oct2025.pdf).
- For the latest information on COVID-19 and prevention measures, please visit the thematic website of COVID-19 (<https://www.coronavirus.gov.hk/eng/index.html>).

Laboratory surveillance for COVID-19 cases

Positive nucleic acid test laboratory detections for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) virus

(Note: The data reported are provisional figures and subject to further revision.)

In week 35, the weekly number of newly recorded positive nucleic acid test laboratory detections for SARS-CoV-2 virus was 91 as compared to 159 in the preceding week. (Figure 1.1)

In the first 4 days of week 36 (Aug 30– Sep 2), the daily number of newly recorded positive nucleic acid test laboratory detections for SARS-CoV-2 virus ranged from 10 to 16.

Since Jan 30, 2023, the cumulative number of positive nucleic acid test laboratory detections was 89,522 (as of Sep 2, 2026).




Figure 1.1 Weekly number of positive nucleic acid test laboratory detections for SARS-CoV-2 virus

Positive detection rate of specimens tested positive for SARS-CoV-2 virus at the Public Health Laboratory Services Branch, Centre for Health Protection

Among the 9,808 respiratory specimens received by the Public Health Laboratory Services Branch (PHLSB) in week 35, 104 (1.06%) were tested positive for SARS-CoV-2 virus as compared to 204 (2.03%) in the preceding week. (Figure 1.2)




Figure 1.2 Percentage of specimens tested positive for SARS-CoV-2 virus at PHLSB

COVID-19 outbreak surveillance

(Note: The data reported are provisional figures and subject to further revision.)

In week 35, 2 COVID-19 outbreaks occurring in schools/institutions were recorded (affecting 11 persons), as compared to 3 outbreaks recorded in the previous week (affecting 12 persons). (Figure 1.3)

In the first 4 days of week 36 (Aug 30–Sep 2), 4 COVID-19 outbreaks occurring in schools/institutions were recorded (affecting 23 persons).




Figure 1.3 COVID-19 outbreaks in schools/institutions

Type of institutions	Week 34	Week 35	First 4 days of week 36 (Aug 30–Sep 2)
Child care centre/ kindergarten (CCC/KG)	0	0	0
Primary school (PS)	0	0	0
Secondary school (SS)	0	0	0
Residential care home for the elderly	1	1	3
Residential care home for persons with disabilities	2	1	1
Others	0	0	0
<i>Total number of outbreaks</i>	3	2	4
<i>Total number of persons affected</i>	12	11	23

Surveillance of severe and fatal COVID-19 cases

(Note: The data reported are provisional figures and subject to further revision.)

In week 35, the weekly number of severe COVID-19 cases including deaths with cause of death preliminarily assessed to be related to COVID-19 was 3 as compared to 6 in the preceding week. (Figure 1.4)




Figure 1.4 Weekly number of severe COVID-19 cases including deaths

Note: Severe and fatal cases are recorded according to their initial reporting dates.

Sewage surveillance of SARS-CoV-2 virus

In week 35, the 7-day geometric mean per capita viral load of SARS-CoV-2 virus from sewage surveillance was around 110,000 copy/L as compared to around 130,000 copy/L in the preceding week. (Figure 1.5)




Figure 1.5 7-day geometric mean per capita viral load of SARS-CoV-2 virus from sewage surveillance since May 1, 2022

Note: The dotted line refers to the temporary sewage sampling suspension for a safety review by the Drainage Services Department.

Acknowledgement

The initiative is funded by the Hong Kong Jockey Club Charities Trust through its "Special Donation on Epidemic Preparedness" to the CHP.

COVID-19 surveillance among sentinel family medicine clinics and sentinel private medical practitioner clinics

In week 35, the average consultation rate for COVID-19 among sentinel family medicine clinics and sentinel private medical practitioner clinics were 5.6 (Figure 1.6) and 4.1 (Figure 1.7) COVID-19 cases per 1,000 consultations, respectively.




Figure 1.6 Average consultation rate of COVID-19 cases in family medicine clinics




Figure 1.7 Average consultation rate of COVID-19 cases in private medical practitioner clinics

Surveillance on SARS-CoV-2 variants

Currently, WHO is monitoring one variant of interest (VOI), which is JN.1, and four VUMs, which are PQ.16.1.1, NB.1.8.1, XFG and BA.3.2. CHP conducts surveillance on SARS-CoV-2 variants from sewage samples. The latest surveillance data (as of Aug 31, 2026) showed that PQ.16.1.1 is the most prevalent variant, comprising 97.6% of all characterised specimens. (Figure 1.8)




Figure 1.8 Estimated proportion of variants among sewage samples

*Including the variant's descendant lineages, except those individually specified elsewhere in the table.

#Those SARS-CoV-2 variants not classified as VOIs/VUMs by WHO at the time of reporting.

Note: XFG is the descendant lineage of JN.1. BA.3.2 is a descendent lineage of the Omicron variant BA.3.

CHP also conducted genetic characterisation on positive respiratory specimens between Aug 19 and Sep 1, 2026. The results showed that PQ.16.1.1 was the most prevalent variant, comprising 100% of all characterised specimens from severe and fatal cases and 82% from non-severe cases.

Global situation of COVID-19 activity

- According to the WHO, global SARS-CoV-2 activity were stable, with variations observed across some regions.
- The COVID-19 activity in some of the neighbouring regions increased, while the activity levels at Canada, the UK and Australia remained low.
 - ◆ In Chinese Mainland (week ending Aug 23, 2026), the percentage of specimens tested positive for SARS-CoV-2 showed a downward trend. Overall activity remained at a moderate level. The predominant variant was NB.1.8.1 recently. In Taiwan region (week ending Aug 29, 2026), the COVID-19 activity remains in a plateau phase, with the number of COVID-19 outpatient and emergency visits showing a gradual decline. The predominant variant was PQ.16.1.1.
 - ◆ In Japan (week ending Aug 23, 2026), the average number of reported COVID-19 cases per sentinel site was 1.18 compared to 1.13 in the preceding week. The predominant variant was BA.3.2.
 - ◆ In South Korea (week ending Aug 22, 2026), the weekly detection rate for SARS-CoV-2 was 11.3% compared to 9.1% in the preceding week. The predominant variant was BA.3.2.
 - ◆ In Singapore (week ending Aug 22, 2026), the positivity rate for COVID-19 among acute respiratory infection (ARI) samples in the community was 4% compared to 2% in the preceding week.
 - ◆ In the United States (week ending Aug 22, 2026), the percent positivity of COVID-19 was 4.6% compared to 4.8% in the preceding week. The predominant variant was XFG.
 - ◆ In Canada (week ending Jul 25, 2026), indicators of COVID-19 activity remained at low level. The percentage of tests positive for COVID-19 was 1.0%, compared to 1.0% in the preceding week. The predominant variant was NB.1.8.1.
 - ◆ In the United Kingdom (week ending Aug 23, 2026), COVID-19 activity remained at low level. COVID-19 PCR positivity in hospital settings was 2.9% compared to 2.5% in the preceding week. The predominant variant was XFG.
 - ◆ In Europe (week ending Aug 23, 2026), SARS-CoV-2 positivity from sentinel specimens was at 11% compared to 4% at the previous week. The predominant variant was XFG.
 - ◆ In Australia (fortnight ending Aug 23, 2026), test positivity for SARS-CoV-2 has slightly increased. The predominant variant was NB.1.8.1.

Sources:

Information have been extracted from the following sources when updates are available: [World Health Organization](#), [Chinese Center for Disease Control and Prevention](#), [Taiwan Centers for Disease Control](#), [Japan Ministry of Health](#), [Korean Disease Control and Prevention Agency](#), [Singapore Communicable Diseases Agency](#), [United States Centers for Disease Control and Prevention](#), [Public Health Agency of Canada](#), [UK Health Security Agency](#), [European Centre for Disease Prevention and Control \(ECDC\)](#) and [WHO Regional Office for Europe \(WHO Euro\)](#), and [Australian Department of Health and Aged Care](#).

Local Situation of Influenza Activity (as of Sep 2, 2026)

Reporting period: Aug 23 – 29, 2026 (Week 35)

- Hong Kong is in influenza season. The latest surveillance data showed that the overall influenza activity increased and was at a high level.
- Influenza can cause serious illnesses in high-risk individuals and even healthy persons. Given that seasonal influenza vaccines are safe and effective, all persons aged 6 months or above except those with known contraindications are recommended to receive influenza vaccine to protect themselves against seasonal influenza and its complications, as well as related hospitalisations and deaths.
- 2026/27 Seasonal Influenza Vaccination (SIV) Programmes, including the SIV School Outreach Programme (SIVSOP), the Residential Care Home Vaccination Programme and the Vaccination Subsidy Scheme (VSS), will commence on September 17. Eligible persons can receive free or subsidised vaccinations through the SIV Programmes. The public may visit the CHP's Vaccination Schemes page for more details of the vaccination programmes (<https://www.chp.gov.hk/en/features/17980.html>).
- Apart from getting influenza vaccination, members of the public should always maintain good personal and environmental hygiene.
- For the latest information on seasonal influenza and its prevention, please visit the Centre for Health Protection's Seasonal Influenza page (http://www.chp.gov.hk/en/view_content/14843.html).

Influenza-like-illness surveillance among sentinel family medicine clinics and sentinel private medical practitioner clinics, 2022-26

In week 35, the average consultation rate for influenza-like illness (ILI) among sentinel family medicine clinics (FMC) was 17.5 ILI cases per 1,000 consultations, which was higher than 14.4 recorded in the previous week (Figure 2.1, left). The average consultation rate for ILI among sentinel private medical practitioner (PMP) clinics was 40.5 ILI cases per 1,000 consultations, which was lower than 40.9 recorded in the previous week (Figure 2.1, right).




Figure 2.1 ILI consultation rates at sentinel FMC (left) and PMP clinics (right), 2022-26

Laboratory surveillance, 2022-26

Among the 10,279 respiratory specimens* received in week 35, 1,148 (11.17%) were tested positive for seasonal influenza A or B viruses. Among the subtyped influenza detections, there were 846 (76%) influenza A(H1), 206 (19%) influenza A(H3) and 60 (5%) influenza B viruses. The positive percentage (11.17%) was above the baseline threshold of 4.94% and was higher than 10.29% recorded in the previous week (Figure 2.2).




Figure 2.2 Percentage of respiratory specimens tested positive for influenza viruses, 2022-26 (upper: overall positive percentage, lower: positive percentage by subtypes)

[Notes: The Centre for Health Protection (CHP) of the Department of Health closely monitors the local seasonal influenza activity through a series of surveillance systems. Among them, the CHP sets threshold levels for two important influenza indicators, including the positive percentage of influenza detections among respiratory specimens and the admission rate of patients diagnosed with influenza in public hospitals. These threshold levels are calculated statistically based on data collected for both indicators in the past years during non-season periods. Using these thresholds, the CHP assesses the current local situation of seasonal influenza with higher accuracy and determines whether Hong Kong enters influenza season. The CHP annually reviews and analyses the latest surveillance data, and updates these threshold levels where appropriate. The sensitivity of the surveillance system is enhanced with the updated thresholds of positive percentage of influenza detection and admission rate of higher coherence.]

Remarks: Some specimens may contain vaccine strains from people with recent history of receiving live-attenuated influenza vaccine

* Including 9,808 specimens received by Public Health Laboratory Services Branch, Centre for Health Protection and 471 specimens received by the Hospital Authority

Surveillance of antiviral susceptibility of influenza A and B viruses

- The Public Health Laboratory Services Branch of the Centre for Health Protection tests virus isolates of influenza viruses obtained from cell culture for reduced antiviral susceptibility. For influenza A(H1) viruses, genotypic assay for H275Y substitution (which confers resistance to oseltamivir) is also performed on selected clinical specimens.
- In July 2026, there were no new reports of influenza viruses with reduced susceptibility to oseltamivir and baloxavir. There were also no reports of influenza A(H1) virus with H275Y substitution.
- For the results of previous months, please refer to the following webpage: <https://www.chp.gov.hk/en/statistics/data/10/641/695/7124.html>
- The detection rates of oseltamivir-resistant influenza A and B viruses remain low (less than 5%) according to latest surveillance data of overseas countries.

Influenza-like illness outbreak surveillance, 2022-26

In week 35, 29 ILI outbreaks occurring in schools/institutions were recorded (affecting 198 persons), as compared to 18 outbreaks recorded in the previous week (affecting 131 persons) (Figure 2.3). The overall number was at the low intensity level currently (Figure 2.4*). In the first 4 days of week 36 (Aug 30 to Sep 2), 25 ILI outbreaks in schools/institutions were recorded (affecting 104 persons). Since the start of the influenza season in week 26, 272 outbreaks were recorded (as of September 2).




Figure 2.3 ILI outbreaks in schools/institutions, 2022-26

Type of institutions	Week 34	Week 35	Cumulative number of outbreaks during influenza season
Child care centre/kindergarten (CCC/KG)	0	5	36
Primary school (PS)	4	4	68
Secondary school (SS)	0	3	12
Residential care home for the elderly	10	12	99
Residential care home for persons with	3	3	27
Others	1	2	30
<i>Total number of outbreaks</i>	18	29	272
<i>Total number of persons affected</i>	131	198	1890

In comparison, 158 to 884 outbreaks were recorded in the same duration of surveillance (10 complete weeks) in the 2023 to 2025 seasons, as compared with 247 outbreaks in the current season (Figure 2.5).




Figure 2.4 ILI outbreaks in schools/institutions, 2025-26




Figure 2.5 Cumulative numbers of ILI outbreaks reported during major influenza seasons, 2023–26

Note: The predominating virus was shown in bracket.

* Various intensity levels applicable for this year were calculated with the moving epidemic method (MEM). For details, please refer to this webpage: https://www.chp.gov.hk/files/pdf/explanatory_note_for_flux_mem_eng.pdf

Influenza-associated hospital admission rates in public hospitals based on discharge coding, 2022-26

In week 35, the overall admission rate in public hospitals with principal diagnosis of influenza was 0.75 (per 10,000 population) as compared to 0.59 recorded in the previous week (Figure 2.6). It was above the baseline threshold of 0.27 but was at the medium intensity level (Figure 2.7*). The influenza-associated admission rates for persons aged 0-5 years, 6-11 years, 12-17 years, 18-49 years, 50-64 years and 65 years or above were 2.90, 1.50, 0.62, 0.19, 0.28 and 1.81 cases (per 10,000 people in the age group) respectively, as compared to 2.58, 0.90, 0.18, 0.15, 0.23 and 1.48 cases in the previous week (Figure 2.6).




Figure 2.6 Influenza-associated hospital admission rates, 2022-26 (upper: overall rate, lower: rates by age groups)

[Notes: The Centre for Health Protection (CHP) of the Department of Health closely monitors the local seasonal influenza activity through a series of surveillance systems. Among them, the CHP sets threshold levels for two important influenza indicators, including the positive percentage of influenza detections among respiratory specimens and the admission rate of patients diagnosed with influenza in public hospitals. These threshold levels are calculated statistically based on data collected for both indicators in the past years during non-season periods. Using these thresholds, the CHP assesses the current local situation of seasonal influenza with higher accuracy and determines whether Hong Kong enters influenza season. The CHP annually reviews and analyses the latest surveillance data, and updates these threshold levels where appropriate. The sensitivity of the surveillance system is enhanced with the updated thresholds of positive percentage of influenza detection and admission rate of higher coherence.]




Figure 2.7 Influenza-associated hospital admission rates, 2025-26

*Various intensity levels applicable for this year were calculated with the moving epidemic method (MEM). For details, please refer to this webpage: https://www.chp.gov.hk/files/pdf/explanatory_note_for_flux_mem_eng.pdf

Rate of ILI syndrome group in accident and emergency departments, 2022-26[#]

In week 35, the rate of the ILI syndrome group in the accident and emergency departments (AEDs) was 135.5 (per 1,000 coded cases), which was lower than the rate of 137.5 in the previous week (Figure 2.8).

#Note: This syndrome group includes codes related to ILI such as influenza, upper respiratory tract infection, fever, cough, throat pain, and pneumonia.




Figure 2.8 Rate of ILI syndrome group in AEDs, 2022-26

Fever surveillance at sentinel child care centres/kindergartens, 2022-26

As most child care centres / kindergartens (CCCs/KGs) were on summer holiday in weeks 34 and 35, sentinel surveillance was suspended during both weeks. In week 33, 0.46% of children in the sentinel CCCs/KGs had fever (38°C or above) (Figure 2.9).




Figure 2.9 Percentage of children with fever at sentinel CCCs/KGs, 2022-26

Fever surveillance at sentinel residential care homes for the elderly, 2022-26

In week 35, 0.22% of residents in the sentinel residential care homes for the elderly (RCHes) had fever (38°C or above), compared to 0.14% recorded in the previous week (Figure 2.10).




Figure 2.10 Percentage of residents with fever at sentinel RCHes, 2022-26

Influenza-like illness surveillance among sentinel Chinese medicine practitioners, 2022-26

In week 35, the average consultation rate for ILI among Chinese medicine practitioners (CMPs) was 1.24 ILI cases per 1,000 consultations as compared to 0.97 recorded in the previous week (Figure 2.11).




Figure 2.11 ILI consultation rate at sentinel CMPs, 2022-26

Surveillance of severe influenza cases

(Note: The data reported are provisional figures and subject to further revision.)

Surveillance for intensive care unit (ICU) admission/death with laboratory confirmation of influenza among adult patients (Aged 18 years or above)

Since 2018, the Centre for Health Protection (CHP) has collaborated with the Hospital Authority and private hospitals to monitor ICU admissions and deaths with laboratory confirmation of influenza among adult patients regularly. For surveillance purposes, the cases refer to laboratory-confirmed influenza patients who required ICU admission or died within the same admission of influenza infection. Their causes of ICU admission or death may be due to other acute medical conditions or underlying diseases.

- In week 35, 66 adult cases of ICU admission/death with laboratory confirmation of influenza (including 40 deaths) were recorded, as compared to 53 cases (including 38 deaths) in the previous week. Among the 66 adult cases, 33 were not known to have received the SIV. In the first 4 days of week 36 (Aug 30 to Sep 2), 26 cases were recorded, in which 16 of them were fatal.

Week	Influenza type					
	A(H1)	A(H3)	A (pending subtype)	B	A and B	C
Week 35	46	15	2	2	0	1
First 4 days of week 36 (Aug 30 to Sep 2)	16	5	4	0	0	1

- During the influenza season, 467 adult cases of ICU admission/death with laboratory confirmation of influenza were recorded, in which 297 of them were fatal (as of Sep 2). Among them, 223 patients had influenza A(H1) infection, 184 patients with influenza A(H3), 28 patients with influenza B, 30 patients with influenza A (subtype pending) and 2 with influenza C.
- In comparison, 308 to 470 adult cases were recorded in the same duration of surveillance (10 complete weeks) in the 2023 to 2025 seasons, as compared with 441 cases in the current season (Figure 2.12, left). The corresponding figures for deaths were 199 to 318 in the above seasons, as compared with 281 deaths in the current season (Figure 2.12, right).





Figure 2.12 Cumulative numbers of adult severe influenza cases reported during major influenza seasons, 2023–26 (left: ICU admission/death cases; right: deaths)

Note: The predominating virus was shown in bracket.

Surveillance of severe paediatric influenza-associated complication/death (Aged below 18 years)

- In week 35 and the first 4 days of week 36 (Aug 30 to Sep 2), there was 1 case of severe paediatric influenza-associated complication/death.

Reporting week	Age	Sex	Complication	Fatal case?	Influenza subtype	History of receiving 2025/26 influenza vaccine
36	5 years	Male	Severe pneumonia	No	Influenza A(H1)	Yes

- During the influenza season, 12 paediatric cases of influenza-associated complication/death were reported, in which one of them was fatal. Six cases had infections with influenza A(H3) and another six with influenza A(H1). Four cases had not received SIV. In 2026, 20 paediatric cases of influenza-associated complication/death were reported, in which one of them was fatal (as of Sep 2).
- In comparison, 10 to 17 paediatric cases of influenza-associated complication/death were recorded in the same duration of surveillance (10 complete weeks) in the 2023 to 2025 seasons, as compared with 11 cases in the current season (Figure 2.13, left). The corresponding figures for deaths were 0 to 2 in the above seasons, as compared with 1 death in the current season (Figure 2.13, right).




Figure 2.13 Cumulative numbers of cases of paediatric influenza-associated complication/death reported during major influenza seasons, 2023–26 (left: complication/death cases; right: deaths)

Note: The predominating virus was shown in bracket.

Severe influenza cases of all ages

- During the influenza season, 479 severe influenza cases among all ages have been reported, including 298 deaths (as of Sep 2).

Age group	Cumulative number of cases (death)
0-5	6 (0)
6-11	4 (1)
12-17	2 (0)
18-49	31 (4)
50-64	56 (19)
>=65	380 (274)

- Among the adult fatal cases with available clinical information, about 85% had chronic diseases.
- Among patients with laboratory confirmation of influenza admitted to public hospitals in this season, 2.7% of admitted cases died during the same episode of admission. It was within the historical range of 2.5% to 4.5%.




Figure 2.14 Weekly number of severe influenza cases by age groups, 2022-26 (the percentage positive for influenza viruses in Figure 2.2 is also shown in this graph)

Sewage surveillance for seasonal influenza

CHP has leveraged established infrastructure to launch a new sewage-based surveillance indicator for tracking local seasonal influenza activity in the community as a complement to the conventional systems. Starting from late October 2025, CHP publishes sewage surveillance results on seasonal influenza viruses.

In week 35, viral load of influenza A and B viruses from sewage surveillance was 5.42 copies (unit adjusted for population)*, which was higher than 3.88 copies* recorded in the previous week (Figure 2.15).




Figure 2.15 Viral load of influenza A and B viruses from sewage surveillance, 2026

**Note: The unit for influenza viral load in sewage is the number of influenza virus copies per 1,000 copies of Pepper Mild Molt Virus (PMMoV), which is a standardised unit. PMMoV is a plant virus primarily found in peppers and their products. It is harmless to humans and does not cause disease. It enters the human body through food ingestion and is excreted in faeces, making it a common biomarker for estimating the size of the population catchment of the sampling sites. Normalising viral load data with PMMoV can reduce the influence of population fluctuations, thereby making the monitoring more accurate and reliable.*

Acknowledgement

The initiative is funded by the Hong Kong Jockey Club Charities Trust through its "Special Donation on Epidemic Preparedness" to the CHP.

Global Situation of Influenza Activity

Globally, influenza detections remained low in week 34, 2026. Influenza A and B viruses were detected in similar proportions (data up to Aug 23, 2026).

- In the United States (week ending Aug 22, 2026), seasonal influenza activity was low. The percentage of specimens tested positive for influenza was below 1%.
- In Europe (week ending Aug 23, 2026), influenza activity remained at a low level. Influenza positivity increased from 4% in the previous week to 6%, but remained below the 10% epidemic threshold.
- In the United Kingdom (week ending Aug 23, 2026), influenza activity increased but remained at baseline levels.
- In Chinese Mainland (week ending Aug 23, 2026), the percentage of specimens that tested positive for influenza in the southern provinces slightly decreased, while that in the northern provinces remained low, with the rates at 18.3% and 2.0% in week 34, respectively. The proportion of influenza A(H3N2) among detections increased nationwide. In Macao (week ending Aug 22, 2026), the influenza positivity rate decreased compared with the previous week. Influenza A (H3) was the predominant virus detected.
- Japan entered the influenza season in late August. In Japan (week ending Aug 23, 2026), influenza activity increased, with the average number of reported ILI cases per sentinel site rising to 1.11 in week 34, exceeding the epidemic baseline threshold of 1.0. Influenza A(H1) was the predominant virus detected.
- In South Korea (week ending Aug 22, 2026), influenza activity increased. The weekly proportion of influenza-like illness increased from 7.5 to 9.2 per 1,000 consultations compared with the previous week. Influenza positivity also increased from 5.4% to 10.0%. Influenza A(H1) was the predominant virus detected.
- In Australia (fortnight ending Aug 23, 2026), the number of influenza cases increased substantially in the last fortnight and current surveillance data suggest that the national influenza peak in 2026 is yet to occur. Majority of the influenza notifications were influenza A.
- In New Zealand (week ending Aug 23, 2026), the national ILI rate increased to 38.29 per 100,000 population, higher than the rate reported to this time in 2022–2025. Among sentinel influenza-positive samples, influenza A(H3) was the most frequently detected subtype (68%), followed by A(H1) (21%).

Sources:

Information has been extracted from the following sources when updates are available: [World Health Organization](#), [United States Centers for Disease Control and Prevention](#), [European Centre for Disease Prevention and Control \(ECDC\)](#) and [WHO Regional Office for Europe \(WHO Euro\)](#), [UK Health Security Agency](#), [Chinese National Influenza Center](#), [Australian Centre for Disease Control](#), [New Zealand Ministry of Health](#), [Japan Ministry of Health](#) and [Korean Disease Control and Prevention Agency](#).