
登革熱 Dengue Fever

由登革熱病毒引起


Caused by Dengue viruses



病徵
Symptoms

高燒、嚴重頭痛、眼窩後疼痛、肌肉及關節痛、噁心、嘔吐、淋巴結腫脹以及出疹


High fever, severe headache, pain behind the eyes, muscle and joint pain, nausea, vomiting, swollen lymph nodes and rash



潛伏期
Incubation period


3–14天，通常為 4–7天

Ranging from 3 – 14 days, commonly 4 – 7 days



外遊返港後若感到不適，應盡快求醫，並將行程細節告知醫生

Anyone feeling unwell after a trip should seek medical advice as soon as possible and provide travel details to their doctors



登革熱病毒
Dengue viruses

- 有四種不同血清型
- 首次感染登革熱的病情一般較輕，若隨後感染其他血清型的病毒，出現重症登革熱(又稱「登革出血熱」)的機會較高
- 縱使病癒，仍終身有風險染上重症登革熱
- 重症登革熱是一種由登革熱所引起的併發症，病情比較嚴重，可導致死亡

- Encompass 4 different serotypes
- The symptoms of first infection are usually mild, but subsequent infections with other serotypes are more likely to result in severe dengue (also known as 'dengue haemorrhagic fever')
- Even though recovered, still have life-long risk of contracting severe dengue
- Severe dengue is a severe and potentially fatal complication of dengue fever



預防方法
Prevention

- 清除積水
- 避免蚊叮

- Eliminate pockets of stagnant water
- Avoid mosquito bites



衛生防護中心 Centre for Health Protection

衛生防護中心網站 Centre for Health Protection Website
www.chp.gov.hk

衛生防護中心 Facebook 專頁 Centre for Health Protection Facebook Fanpage
fb.com/CentreforHealthProtection

衛生署健康教育專線 Health Education Infoline of the Department of Health
2833 0111

衛生署 Department of Health

Revised in July 2023
二零二三年七月修訂