

# FLU EXPRESS



*Flu Express* is a weekly report produced by the Respiratory Disease Office of the Centre for Health Protection. It monitors and summarizes the latest local and global influenza activities.

## Local Situation of Influenza Activity (as of Aug 29, 2018)

**Reporting period: August 19 – 25, 2018 (Week 34)**

- The latest surveillance data showed that the overall influenza activity in Hong Kong remained at a low level.
- Influenza can cause serious illnesses in high-risk individuals and even healthy persons. Given that seasonal influenza vaccines are safe and effective, all persons aged 6 months or above except those with known contraindications are recommended to receive influenza vaccine to protect themselves against seasonal influenza and its complications, as well as related hospitalisations and deaths.
- In the coming 2018/19 season, the Vaccination Subsidy Scheme (VSS) will be expanded to cover those aged 50 to 64 to receive subsidised seasonal influenza vaccination. It will also continue to provide subsidised vaccination to children aged 6 months to under 12 years, elderly aged 65 years or above, pregnant women, persons with intellectual disabilities and recipients of Disability Allowance. In the Government Vaccination Programme (GVP), eligible groups for free vaccination will be the same as that of 2017/18. VSS and GVP will be launched on 10 and 24 October respectively. For more details, please refer to the webpage ([http://www.chp.gov.hk/en/view\\_content/17980.html](http://www.chp.gov.hk/en/view_content/17980.html)).

## Influenza-like-illness surveillance among sentinel general outpatient clinics and sentinel private doctors, 2014-18

In week 34, the average consultation rate for influenza-like illness (ILI) among sentinel general outpatient clinics (GOPCs) was 3.1 ILI cases per 1,000 consultations, which was higher than 2.8 recorded in the previous week (Figure 1, left). The average consultation rate for ILI among sentinel private doctors was 44.2 ILI cases per 1,000 consultations, which was higher than 36.4 recorded in the previous week (Figure 1, right).




Figure 1 ILI consultation rate at sentinel GOPCs (left) and private doctors (right), 2014-18

## Laboratory surveillance, 2014-18

Among the respiratory specimens received in week 34, the positive percentage of seasonal influenza viruses was 2.46%, which was below the baseline threshold of 10.7% and was lower than 2.96% recorded in the previous week (Figure 2). The 103 influenza viruses detected last week included 77 (1.84%) influenza A(H1), 24 (0.57%) influenza A(H3), 1 (0.02%) influenza B and 1 (0.02%) influenza C.




Figure 2 Percentage of respiratory specimens tested positive for influenza viruses, 2014-18 (upper: overall positive percentage, lower: positive percentage by subtypes)

[Note: The baseline threshold is 1.96 standard deviation above the average weekly positive percentage during non-season periods from 2014-2017.]

## Influenza-like illness outbreak surveillance, 2014-18

In week 34, two ILI outbreaks occurring in schools/institutions were recorded (affecting 17 persons), as compared to three outbreaks recorded in the previous week (affecting 14 persons) (Figure 3). In the first 4 days of week 35 (Aug 26 to 29), no ILI outbreaks in schools/institutions were recorded.




Figure 3 ILI outbreaks in schools/institutions, 2014-18

Type of institutions	Week 33	Week 34	First 4 days of week 35 (Aug 26 to 29)
Kindergarten/ child care centre	1	1	0
Primary school	1	0	0
Secondary school	0	0	0
Residential care home for the elderly	1	1	0
Residential care home for persons with disabilities	0	0	0
Others	0	0	0
<i>Total number of outbreaks</i>	3	2	0
<i>Total number of persons affected</i>	14	17	0

## Influenza-associated hospital admission rates in public hospitals based on discharge coding, 2014-18

In week 34, the overall admission rates in public hospitals with principal diagnosis of influenza was 0.08 (per 10,000 population), which was below the baseline threshold of 0.20 but was the same as that recorded in the previous week. The influenza-associated admission rates for persons aged 0-4 years, 5-9 years, 10-64 years and 65 years or above were 0.76, 0.26, 0.03 and 0.12 cases (per 10,000 people in the age group) respectively, as compared to 0.51, 0.49, 0.03 and 0.13 cases in the previous week (Figure 4).




Figure 4 Influenza-associated hospital admission rates, 2014-18 (upper: overall rate, lower: rates by age groups)

[Note: The baseline threshold is 1.96 standard deviation above the average weekly admission rate during non-season periods from 2014-2017.]

## Rate of ILI syndrome group in accident and emergency departments, 2014-18<sup>#</sup>

In week 34, the rate of the ILI syndrome group in the accident and emergency departments (AEDs) was 130.8 (per 1,000 coded cases), which was higher than the rate of 122.5 in the previous week (Figure 5).

*#Note: This syndrome group includes codes related to ILI such as influenza, upper respiratory tract infection, fever, cough, throat pain, and pneumonia.*




Figure 5 Rate of ILI syndrome group in AEDs, 2014-18

## Fever surveillance at sentinel child care centres/ kindergartens, 2014-18

The surveillance for week 34 was suspended due to summer holiday. In week 33, 0.69% of children in the sentinel child care centres / kindergartens (CCCs/KGs) had fever (38°C or above) as compared to 0.67% recorded in the previous week (Figure 6).




Figure 6 Percentage of children with fever at sentinel CCCs/KGs, 2014-18

## Fever surveillance at sentinel residential care homes for the elderly, 2014-18

In week 34, 0.09% of residents in the sentinel residential care homes for the elderly (RCHes) had fever (38°C or above), compared to 0.12% recorded in the previous week (Figure 7).




Figure 7 Percentage of residents with fever at sentinel RCHes, 2014-18

## Influenza-like illness surveillance among sentinel Chinese medicine practitioners, 2014-18

In week 34, the average consultation rate for ILI among Chinese medicine practitioners (CMPs) was 2.07 ILI cases per 1,000 consultations as compared to 1.49 recorded in the previous week (Figure 8).




Figure 8 ILI consultation rate at sentinel CMPs, 2014-18

## Surveillance of severe influenza cases

(Note: The data reported are provisional figures and subject to further revision.)

### **Surveillance for intensive care unit (ICU) admissions/deaths with laboratory confirmation of influenza among adult patients (Aged 18 years or above)**

*For surveillance purpose, the cases refer to laboratory-confirmed influenza patients who required ICU admission or died within the same admission of influenza infection. Their causes of ICU admission or death may be due to other acute medical conditions or underlying diseases.*

- In week 34, four adult cases of ICU admission/death with laboratory confirmation of influenza (including four deaths) were recorded while no cases were recorded in the previous week. Two of the four severe adult cases were known to have received the 2017/18 influenza vaccine.

Week	Influenza type				
	A(H1)	A(H3)	B	C	A (pending subtype)
Week 33	0	0	0	0	0
Week 34	1	1	0	0	2

### **Surveillance of severe paediatric influenza-associated complication/death (Aged below 18 years)**

- In week 34 and the first 4 days of week 35 (Aug 26 to 29), there was one case of severe paediatric influenza-associated complication/death.

Reporting week	Age	Sex	Complication	Fatal case?	Influenza subtype	History of receiving influenza vaccine for this season
34	6 years	Male	Encephalopathy	No	Influenza A(H1)	Yes

Data as of Aug 29, 2018

- In 2018, 26 paediatric cases of influenza-associated complication/death were recorded, in which three of them were fatal (as of Aug 29). 21 (81%) did not receive the influenza vaccine for the 2017/18 season.




Figure 9 Weekly number of severe influenza cases by age groups, 2014-18 (the percentage positive for influenza viruses in Figure 2 is also shown in this graph)

Note: The surveillance system for severe influenza cases among adult patients aged 18 years or above was only activated intermittently during influenza seasons before 2018.



## Surveillance of oseltamivir resistant influenza A(H1N1)pdm09 virus infection

- In week 34 and the first 4 days of week 35 (Aug 26 to 29), there were no new reports of oseltamivir (Tamiflu) resistant influenza A(H1N1)pdm09 virus infection. There are totally 48 reports of oseltamivir resistant influenza A(H1N1)pdm09 virus detected in Hong Kong since 2009.

## Global Situation of Influenza Activity

Influenza activity remained at inter-seasonal levels in most of the temperate zone of the northern hemisphere. Reports of influenza activity were reported as decreased in some countries of tropical America. In the temperate zones of southern hemisphere, influenza activity remained elevated in South America and continued to decrease in Southern Africa. Influenza activity remained below seasonal threshold in Australia and New Zealand. Worldwide, seasonal influenza subtype A viruses accounted for the majority of detections.

- In the United States (week ending Aug 18, 2018), influenza activity was at a low level. The proportion of outpatient visits for ILI was 0.8%, which was below the national baseline of 2.2%. The percentage of respiratory specimens testing positive for influenza was 0.32%.
- In the United Kingdom (week ending Aug 12, 2018), indicators for influenza showed low levels of activity. The positivity of influenza detection was 0.7% in the week ending August 12, 2018, which was below the baseline threshold of 8.6%.
- In Australia (fortnight ending Aug 12, 2018), influenza activity was low and remained within or below the bounds of previous years. Majority of confirmed influenza cases reported nationally were influenza A (87%).
- In New Zealand (week ending August 26, 2018), influenza activity was still unseasonably low. Influenza detection increased in GP visits but decreased in the hospital last week. Influenza A(H1N1) still predominates where detected this season.
- In Mainland China (week ending Aug 19, 2018), influenza activity remained at an extremely low level. There were only few detections of influenza viruses in southern provinces, which were mainly influenza A(H1N1) viruses .
- In Macau (week ending Aug 11, 2018), the proportions of ILI cases in emergency departments among adults was stable and that among children increased as compared to the previous week. The proportion of influenza detections also increased from the previous week.

### Sources:

Information have been extracted from the following sources when updates are available: [World Health Organization](#), [United States Centers for Disease Control and Prevention](#), [Public Health England](#), [Australian Department of Health](#), [New Zealand Ministry of Health](#), [Chinese National Influenza Center](#) and [Health Bureau of Macao Special Administrative Region](#).