

What is melioidosis?

Melioidosis is a disease caused by a bacterium *Burkholderia pseudomallei* which is widespread in soils and muddy water. It can be transmitted through contact with contaminated soil and surface waters (especially through skin abrasions/wounds), inhalation of contaminated dust/water droplets and ingestion of contaminated water. Person-to-person transmission is rare but may occur through contact with the blood or body fluids of an infected person. *Burkholderia pseudomallei* in the soil and muddy water may be exposed to the ground after typhoons or storms, and the bacteria would spread more easily with strong wind and storms. As such, infection cases are more common after typhoons or storms. Depending on the site of infection, common symptoms include fever, headache, localised pain or swelling, ulceration, chest pain, cough, haemoptysis, and swelling of regional lymph nodes.



High risk groups

Persons with underlying diseases including diabetes, lung disease, liver disease, renal disease, cancer, or immunosuppression have higher risks for contracting the disease.

Prevention

- ✓ Avoid contact with contaminated soil.
- ✓ Wear appropriate protective clothing or footwear when participating in activities with possible contact with soil or water, e.g. use gloves and wear boots. High risk individuals may consider to wear a surgical mask in addition.
- ✓ Wash or shower after exposure to contaminated water or soil. Let tap water run for one minute at the start of the day before using it for brushing teeth, washing face or bath, in particular for high risk individuals.
- ✓ Always clean any wounds as soon as possible and cover any cuts or grazes with waterproof dressings.
- ✓ Wash hands with liquid soap and water after handling with soils or gardening.
- ✓ Melioidosis can be spread by contaminated soil and water during and after typhoons and storms. Whether practicable, stay indoor during typhoons and storms, avoid travelling to areas with potential flooding, and not to wade or contact with muddy water and soil. In addition, high risk individuals should avoid path near stormwater drains where aerosols may be generated from contaminated water. If possible exposure is inevitable, high-risk individuals should wear a surgical mask, gloves and boots.
- ✓ Observe food hygiene and avoid drinking unboiled or untreated water.
- ✓ Travellers can contract the disease through outdoor water sports. Risk of infection can be minimised by avoiding exposure to water sources (such as rivers, ponds or lakes) that might be contaminated.

Please scan the following QR codes for updated information:



繁體中文



English



Hindi
हिन्दी



Nepali
नेपाली



Urdu
اردو



Thai
ไทย



Bahasa Indonesia



Tagalog



Sinhala
සිංහල භාෂාව



Bengali
বাংলা ভাষা



Vietnamese
Tiếng Việt

