

寨卡病毒感染 Zika Virus Infection

由寨卡病毒引起的蚊媒傳染病

Mosquito-borne disease caused by Zika virus




病徵 Symptoms

皮疹、發燒、結膜炎、肌肉和關節疼痛、疲累和頭痛。
病徵輕微和持續2-7天

Skin rash, fever
conjunctivitis, muscle and joint pain, malaise and headache.

Symptoms are mild and last for 2-7 days



潛伏期 Incubation period


寨卡病毒感染的潛伏期為3-14天

The incubation period of Zika virus infection ranges from 3–14 days



外遊返港後若感到不適，應盡快求診，並將行程細節告知醫生

Seek medical advice promptly and provide travel details to doctor if feeling unwell after a trip



預防方法 Prevention

- 避免蚊叮
- 防止蚊蟲滋長
- 採取措施以預防性接觸傳染寨卡病毒
- Avoid mosquito bites
- Help prevent mosquito proliferation
- Take precautions to prevent sexual transmission of Zika virus



衛生防護中心
Centre for Health Protection



衛生防護中心網站
Centre for Health Protection Website
www.chp.gov.hk



衛生防護中心 Facebook 專頁
Centre for Health Protection Facebook
fb.com/CentreforHealthProtection

衛生署健康教育專線
Health Education Infoline of the Department of Health

2833 0111



衛生署
Department of Health
二零二三年六月修訂
Revised in June 2023