

Recommended PPE for routine patient care and performing aerosol-generating procedures ^(a) in hospitals/clinics for suspected or confirmed cases with Novel Influenza infection under different response levels

Apply standard precautions (SP) +/- transmission-based precautions for all patients			
Areas	Activities	Recommended PPE	
		Alert/ Serious response level (S1)	Serious response level (S2)/ Emergency response level
High-risk patient areas <ul style="list-style-type: none"> ● Triage stations of Out-patient Clinics and AEDs; ● Designed clinics and ● Isolation rooms(including isolation rooms in ICU and AEDS) 	Routine patient care and aerosol-generating procedures ^(a, b)	<ul style="list-style-type: none"> ● N95 respirator ● eye protection ^(e) ● gown ● gloves ● cap(optional) <p><i>Surgical mask could be an alternative for triage stations based on nature of encounter upon risk assessment</i></p>	
	No patient contact (eg. Outside patient room)	<ul style="list-style-type: none"> ● Surgical mask 	
Other patient areas	Routine patient care	<ul style="list-style-type: none"> ● Standard Precautions +/- transmission based precautions 	<ul style="list-style-type: none"> ● Surgical mask, Standard Precautions +/- transmission based precautions
	aerosol-generating procedures ^(a, c)	<ul style="list-style-type: none"> ● Surgical mask / N95 respirator ^(d) ● eye protection ^(e) ● gown ● gloves 	<ul style="list-style-type: none"> ● N95 respirator ● eye protection ^(e) ● gown ● gloves ● cap (optional).
	No patient contact	<ul style="list-style-type: none"> ● Surgical mask for signs and symptoms of respiratory infection 	<ul style="list-style-type: none"> ● Serious S2 Level: Surgical mask is required in patient areas ● Emergency Level: Surgical mask is required in ALL areas
Non patient areas	Non-patient areas	<ul style="list-style-type: none"> ● Surgical mask for signs and symptoms of respiratory infection 	<ul style="list-style-type: none"> ● Surgical mask is required in patient areas ● Emergency Level: Surgical mask is required in ALL areas

Remarks

- a. Aerosol-generating procedures with documented increased in risk of respiratory infection transmission are endotracheal intubation, cardiopulmonary resuscitation, bronchoscopy, open suctioning of respiratory tract (including tracheostomy care), autopsy and non-invasive positive pressure ventilation (BiPAP & CPAP). Aerosol-generating procedures with controversial/ limited studies evaluating the risk of respiratory infection transmission are high-frequency oscillatory ventilation, nebulizer therapy and sputum induction. Nasopharyngeal aspiration (NPA) and high flow oxygen are theoretically at risk of dispersal of infectious respiratory droplets, therefore they should be performed in conditions as required for aerosol-generating procedures in high-risk patient areas. Other procedures should be assessed on discretion of hospital Infection Control Officers.
- b. In high risk patient areas, place patient in a negative pressure airborne infection isolation room (AIIR) before performing aerosol generating procedures.
- c. In other patient areas, place patient in a well-ventilated area (e.g .at least minimum overall 6 air changes per hour (ACH) or use portable HEPA filter e.g. IQ Air if indicated) before performing aerosol generating procedures.
- d. Based on transmission-based precautions and risk assessment
- e. Eye protection refers to full face shield or goggles or eye-visors.

Centre for Health Protection

Dec 2013

(Last updated on 11 July 2016)

Infection Control Branch, Centre for Health Protection