

Seasonal Influenza Vaccination

Information on Side Effects (Injectable Vaccine)

and 2nd dose Arrangement

The Department of Health (DH) has arranged Vaccination Team (by DH or through public private partnership) to provide your child _____ (name of student) with Seasonal Influenza Vaccine (SIV) at your child's school on _____ (date). Inactivated SIV (by injection) was provided. Please note the information below:

1. Inactivated influenza vaccine is very safe and usually well tolerated, apart from occasional soreness, redness or swelling at the vaccination site.
2. Some children may experience fever, muscle pain, and tiredness 6 to 12 hours after vaccination. These usually improve in two days.
3. If fever or discomfort persists, please consult a doctor. Severe allergic reactions like hives, swelling of the lips or tongue, and difficulties in breathing, or serious adverse events such as limb numbness or weakness are rare but require emergency consultation.

The Vaccination Team will visit the school again on _____ to provide 2nd dose vaccination for your child. (Children under 9 years old who have never received any SIV are recommended to have 2 doses of SIV with a minimum interval of 4 weeks.)

If you have any queries regarding SIV, please call _____

Vaccination Team from: _____

(Name of Enrolled doctor/ Medical Organisation)

季節性流感疫苗 副作用資料頁 (注射式疫苗)及第二劑的安排

衛生署已於 _____ (日期) 安排疫苗接種隊 (由衛生署或透過公私營合作) 到校為 貴子女 _____ (學生姓名) 接種滅活季節性流感疫苗 (注射式)。請留意以下資訊：

1. 滅活流感疫苗十分安全，除了接種部位可能會出現痛楚、紅腫外，一般並無其他副作用。
2. 部分學生在接種後 6 至 12 小時內可能會出現發燒、肌肉疼痛，以及疲倦等症狀，這些症狀通常會在兩天內減退。
3. 如持續發燒或不適，請諮詢醫生意見。若出現罕見的風疹塊、口舌腫脹及呼吸困難等嚴重過敏反應，或嚴重不良情況，如：手腳麻痺、無力，患者必須立即求醫。

接種隊將於 _____ (日期) 再次到校為 貴子女接種第二劑疫苗。(9 歲以下從未接種過季節性流感疫苗的學生，建議應接種兩劑流感疫苗，而兩劑疫苗的接種時間必須最少相隔 4 星期。)

如有任何關於季節性流感疫苗的疑問，請致電

接種隊: _____

(已配對醫生姓名/ 醫療機構名稱)