

## **Seasonal Influenza Vaccination Information on Side Effects (Nasal Spray Vaccine) and 2<sup>nd</sup> dose Arrangement**

---

The Department of Health (DH) has arranged Vaccination Team (by DH or through public private partnership) to provide your child \_\_\_\_\_ (name of student) with Seasonal Influenza Vaccine (SIV) at your child's school on \_\_\_\_\_ (date). Live attenuated SIV (by nasal spray) was provided. Please note the information below:

1. The most common side effects following live attenuated influenza vaccination are fever, nasal congestion or runny nose.
2. If fever or discomfort persists, please consult a doctor. Severe allergic reactions like hives, swelling of the lips or tongue, and difficulties in breathing are rare but require emergency consultation.

The Vaccination Team will visit the school again on \_\_\_\_\_ to provide 2<sup>nd</sup> dose vaccination for your child. (Children under 9 years old who have never received any SIV are recommended to have 2 doses of SIV with a minimum interval of 4 weeks.)

If you have any queries regarding SIV, please call  
\_\_\_\_\_

Vaccination Team from: \_\_\_\_\_

(Name of Enrolled doctor/ Medical Organisation)

# 季節性流感疫苗

## 副作用資料頁 (噴鼻式疫苗)及第二劑的安排

---

衛生署已於 \_\_\_\_\_ (日期) 安排疫苗接種隊 (由衛生署或透過公私營合作) 到校為 貴子女 \_\_\_\_\_ (學生姓名) 接種減活季節性流感疫苗 (噴鼻式)。請留意以下資訊：

1. 接種減活流感疫苗最常見的副作用包括：發燒、鼻塞或流鼻水。
2. 如持續發燒或不適，請諮詢醫生意見。若出現罕見的風疹塊、口舌腫脹及呼吸困難等嚴重過敏反應，患者必須立即求醫。

接種隊將於 \_\_\_\_\_ (日期) 再次到校為 貴子女接種第二劑疫苗。(9 歲以下從未接種過季節性流感疫苗的學生，建議應接種兩劑流感疫苗，而兩劑疫苗的接種時間必須最少相隔 4 星期。)

如有任何關於季節性流感疫苗的疑問，請致電

---

接種隊: \_\_\_\_\_  
(已配對醫生姓名/ 醫療機構名稱)